

Name: _____

Period: _____

Seat#: _____

Directions: Show all work in a way that would earn you credit on the AP Test! This is always the rule! Some answers are provided at the end in italics and underlined. If you need more space, use binder paper and staple to your worksheet.


Average Bond Enthalpies (kJ/mol)

Single Bonds

C—H	413	N—H	391	O—H	463	F—F	155
C—C	348	N—N	163	O—O	146	Cl—F	253
C—N	293	N—O	201	O—F	190	Cl—Cl	242
C—O	358	N—F	272	O—Cl	203	Br—F	237
C—F	485	N—Cl	200	O—I	234	Br—Cl	218
C—Cl	328	N—Br	243	S—H	339	Br—Br	193
C—Br	276	H—H	436	S—F	327	I—Cl	208
C—I	240	H—F	567	S—Cl	253	I—Br	175
C—S	259	H—Cl	431	S—Br	218	I—I	151
Si—H	323	H—Br	366	S—S	266		
Si—Si	226	H—I	299				
Si—C	301						
Si—O	368						

Multiple Bonds



C=C	614	N=N	418	O ₂	495
C≡C	839	N≡N	941	S=O	523
C=N	615			S=S	418
C≡N	891				
C=O	799				
C≡O	1072				



1) Considering bonds broken and formed ONLY, what is the enthalpy change for the following reaction:




Hint C₁₂H₂₄ is a weird molecule. It looks like this, with all single bonds




2) The rxn $\text{BBr}_3(\text{g}) + \text{BCl}_3(\text{g}) \rightarrow \text{BBr}_2\text{Cl}(\text{g}) + \text{BCl}_2\text{Br}(\text{g})$ has a ΔH very close to zero. Explain why ΔH is so small.

Dougherty Valley HS Chemistry - AP
Bonding – Bond Energy Practice 2


3) Determine the enthalpy of reaction for the following using bond energies.



4) Ammonia reacts with oxygen to form nitrogen dioxide and steam, as follows. Use data for bond energies to determine the bond energy of the N–O bond of NO_2 (Hint N-O bond has resonance. Just do the AVERAGE N-O, assume all bonds are equal)



5) Determine the enthalpy of the following reaction using bond energies:



6) Determine the enthalpy for the following reaction: $\text{C}(\text{s}) + \text{CO}_2(\text{g}) \rightarrow 2\text{CO}(\text{g})$

The enthalpy of sublimation of
graphite, $\text{C}(\text{s})$ is 719 kJ/mol

Dougherty Valley HS Chemistry - AP
Bonding – Bond Energy Practice 2

7) Calculate the bond dissociation energy for one mole of O–F bonds, given the following data. (Hint: oxygen is the central atom of OF₂)



8) Using bond enthalpy (in kJ mol⁻¹) values, determine the heat of formation of methane:

Sublimation energy of
C (s, gr) = 719 kJ/mol

9) An unknown gas, X₂, which behaves much like nitrogen gas (N≡N), is analyzed and the following enthalpies of formation are obtained. The X–H bond energy is known to be 383 kJ/mol. Use this information to estimate the X–X single-bond energy in the X₂H₄ molecule. (Hint Write a rxn forming X₂H₄ to start)

X(g) = 412 kJ/mol
H(g) = 217 kJ/mol
X₂H₄(g) = 3 kJ/mol

Answers (Remember that answers for this topic vary greatly if not using the same bond enthalpy data charts! Not a big deal. Use my numbers when given so that your answers will match)

- 1) 0 KJ/mol
- 2) Breaking and making same number of same type of bonds
- 3) -243 KJ/mol
- 4) 467 KJ/mol
- 5) -1905 KJ/mol
- 6) 173 KJ/mol
- 7) 187 KJ/mol
- 8) -61 KJ/mol
- 9) 157 KJ/mol