

N9 – HALF LIFE



Half-Life


- Half Life is the time required for half of a radioisotope's nuclei to decay into its products.

# of $\frac{1}{2}$ lives	% Remaining
0	100%
1	50%
2	25%
3	12.5%
4	6.25%
5	3.125%
6	1.5625%

Graphing Half-Life

Half of the radioactive atoms decay each half-life.

Decay of Radon-220



Using a Table

- Suppose you have 10.0 grams of strontium – 90, which has a half life of 29 years. How much will be remaining after x number of years?

# of $\frac{1}{2}$ lives	Time (Years)	Amount Remaining (g)
0	0	10
1	29	5
2	58	2.5
3	87	1.25
4	116	0.625

Half-Life Equation

- Use a handy dandy equation!

Amount Starting

$$A_E = A_S \times (0.5)^n$$

of half-lives

Amount Ending

$$n = \frac{t}{h}$$

t = time passed
h = length of one half-life

Solving for % remaining

$$A_E = A_S \times (0.5)^n$$

$$\% \text{ remaining} = \frac{A_E}{A_S} \times 100$$

$$\frac{A_E}{A_S} = (0.5)^n$$

Then multiply your answer by 100 to put it in % format!

Example 1

- If gallium – 68 has a half-life of 68.3 minutes, how much of a 160.0 mg sample is left after 1 half life? 80 mg. After 2 half lives? 40 mg. After 3 half lives? 20 mg

$$A_E = A_s \times (0.5)^n$$

$$80 \text{ mg} = 160.0 \text{ mg} \times (0.5)^1$$

$$40 \text{ mg} = 160.0 \text{ mg} \times (0.5)^2$$

Example 2

Cobalt – 60, with a half-life of 5 years, is used in cancer radiation treatments. If a hospital purchases a supply of 30.0 g, how much would be left after 15 years? 3.75 g

$$A_E = A_s \times (0.5)^n$$

$$A_E = 30.0\text{g} \times (0.5)^{(15/5)} = 3.75 \text{ g}$$

Example 3

Iron-59 is used in medicine to diagnose blood circulation disorders. The half-life of iron-59 is 44.5 days. How much of a 2.000 mg sample will remain after 133.5 days? _____

$$A_E = A_s \times (0.5)^{t/h}$$

$$A_E = 2.000 \text{ mg} \times (0.5)^{(133.5/44.5)}$$

$$0.2500 \text{ mg} = 2.000 \text{ mg} \times 0.125$$

Solve for Time/Half-life

$$A_E = A_s \times (0.5)^{t/h}$$

Isolate $(0.5)^{t/h}$

$$\frac{A_E}{A_s} = (0.5)^{t/h}$$

Bring down exponent using logs

$$\text{Log} \left(\frac{A_E}{A_s} \right) = \frac{t}{h} \text{Log} (0.5)$$

*Plug in your #'s
then rearrange
for t or h
depending on
what you want
to solve for!*

Example 4

The half-life of polonium-218 is 3.0 min. If you start with 20.0 g, how long before only 1.25 g remains?

$$\text{Log} \left(\frac{A_E}{A_S} \right) = \frac{t}{h} \text{Log} (0.5)$$

$$\text{Log} \left(\frac{1.25}{20.0} \right) = \frac{t}{3} \text{Log} (0.5)$$

$$t = 12\text{min}$$

Example 5

A sample initially contains 150.0 mg of radon-222. After 11.4 days, it contains 18.75 mg of radon-222. Calculate the half-life.

$$\text{Log} \left(\frac{A_E}{A_S} \right) = \frac{t}{h} \text{Log} (0.5)$$

$$\text{Log} \left(\frac{18.75}{150.0} \right) = \frac{11.4}{h} \text{Log} (0.5)$$

$$h = 3.8 \text{ days}$$